


Please amend claims 1, 2, 8-11, 16-19, 22 and 44-47. Please cancel claims 12 and 13.


In the claims

1. (currently amended) A compound of formula (1)



wherein

the group R_1-A is a purine radical of formula (2)



X is oxygen;

R_1 is a group $-R_2-L_2$ ~~or a group R_5~~ ;

R_2 and R_4 are, independently of each other, a straight or branched chain alkylene group or polyvalent branched chain alkyl group with 1 to 300 carbon atoms, optionally substituted by a lower alkyl, lower alkoxy, lower acycloxy or halogen wherein optionally

(a) one or more carbon atoms are replaced by oxygen;

(b) one or more carbon atoms are replaced by nitrogen carrying a hydrogen atom, and the adjacent carbon atom is substituted by oxo;

(c) one or more carbon atoms are replaced by oxygen, and the adjacent carbon atom is substituted by oxo;

(d) the bond between two adjacent carbon atoms is a double or a triple bond;

(e) one or more carbon atoms are replaced by a phenylene, a saturated or unsaturated cycloalkylene, a saturated or unsaturated bicycloalkylene, a ~~bridging~~ divalent heteroaromatic or a ~~bridging~~ divalent saturated or unsaturated heterocyclyl group;

(f) two adjacent carbon atoms are replaced by a disulfide linkage; or a combination of two or more alkylene and/or modified alkylene groups as described in ~~defined under~~ (a) to (f) ~~above~~, herein ~~optionally containing substituents~~;

R₃ is an aromatic or a heteroaromatic group, or an optionally substituted 1-alkenylene, 1-alkynylene, 1-cycloalkenylene, or an unsaturated heterocyclyl group with the double bond connected to CH₂;

~~R₅ is an optionally substituted cycloalkyl, cycloalkenyl or heterocyclyl group;~~

R₆ is hydrogen, hydroxy or unsubstituted or substituted amino; one of R₇ ~~and~~ or R₈ is R₁ and the other one is hydrogen; and

L₁ and L₂ are ~~one or a plurality of~~ the same or different labels and each is selected from the group consisting of a spectroscopic probe including a fluorophore or a chromophore, a magnetic probe, a contrast reagent, a radioactive moiety, avidin, streptavidin, biotin, a moiety which is capable of crosslinking to other molecules selected from ~~the group consisting of~~ a maleimide, an active ester, an azide and a benzophenone, a tethered metal-chelate which is capable of generating hydroxyl radicals upon exposure to H₂O₂, ~~and~~ ascorbate, malachite green, a moiety covalently attached to a solid support, a lipid, methotrexate, a linear poly(arginine) of D- and/or L-arginine with 6-15 arginine residues, ~~a linear polymer of 6-15 subunits each carrying a guanidinium group~~, oligomers ~~or short length polymers of~~

6-50 subunits wherein at least one subunit has an attached guanidinium group ~~a portion of which have attached guanidinium groups,~~ and ~~parts~~ a partial amino acid sequence of a sequence of a ~~the~~ HIV-tat protein; or

L_1 is a bond connecting R_4 to A forming a cyclic substrate~~;~~ a further group $-R_3-CH_2-X-A-R_{1,7}$ or a nucleic acid or a derivative thereof capable of undergoing base-pairing with its complementary strand; or L_2 is a nucleic acid or a derivative thereof capable of undergoing base-pairing with its complementary strand if R_7 is hydrogen.

2. (currently amended) The compound according to claim 1, wherein R_3 is phenylene, an unsubstituted or substituted mono- or bicyclic ~~bridging~~ divalent heteroaryl group of 5 or 6 rings atoms comprising zero, one, two, three or four ring nitrogen atoms and zero or one oxygen atom and zero or one sulfur atom, with the proviso that at least one ring carbon atom is replaced by a nitrogen, oxygen or sulfur atom, 1-alkenylene, 1-alkynylene, 1-cyclohexenylene with 3 to 7 carbon atoms, wherein the double or triple bond is connected to CH_2 , or an optionally substituted unsaturated ~~bridging~~ divalent heterocyclyl group with 3 to 12 atoms and 1 to 5 heteroatoms selected from ~~the~~ group consisting of nitrogen, oxygen and sulfur, and a double bond in the position connecting the heterocyclyl group to CH_2 ~~;~~ and ~~R_5 is optionally substituted cycloalkyl with 3 to 7 carbon atoms; optionally substituted cycloalkenyl with 5 to 7 carbon atoms; or optionally substituted saturated or unsaturated heterocyclyl with 3 to 12 atoms, and 1 to 5 heteroatoms selected from the group consisting of nitrogen, oxygen and sulfur.~~

3. (cancelled)

4. (previously presented) The compound according to claim 1, wherein R₃ is phenylene.
5. (previously presented) The compound according to claim 1, wherein R₃ is thienylene.
6. (previously presented) The compound according to claim 1, wherein R₆ is unsubstituted amino, R₇ is R₁, and R₈ is hydrogen.
7. (previously presented) The compound according to claim 1, wherein R₆ is unsubstituted amino, R₇ is a group -R₂-L₂, and R₈ is hydrogen.
8. (currently amended) The compound according to claim 7, wherein L₂ is a ~~spectroscopic probe~~ fluorophore or a chromophore.
9. (currently amended) The compound according to claim 7, wherein ~~L₁ and L₂ are spectroscopic probes~~ is a fluorophore or a chromophore and L₂ is a fluorophore or a chromophore.
10. (currently amended) The compound according to claim 9, wherein each of L₁ and L₂ represents a fluorescence donor/ or a fluorescence quencher ~~pair~~.
11. (currently amended) The compound according to claim 10, wherein each of L₁ and L₂ represents a donor or an acceptor of a FRET pair.
- 12-13. (cancelled)

14. (previously presented) The compound according to claim 1, wherein R_6 is unsubstituted amino, R_7 is hydrogen, and R_8 is R_1 .

15. (previously presented) The compound according to claim 1, wherein R_6 is unsubstituted amino, R_7 is hydrogen, and R_8 is a group $-R_2-L_2$.

16. (currently amended) The compound according to claim 15, wherein L_2 is a ~~spectroscopic probe~~ fluorophore or a chromophore.

17. (currently amended) The compound according to claim ~~15~~ 16, wherein L_1 ~~and L_2 are spectroscopic probes~~ is a fluorophore or a chromophore.

18. (currently amended) The compound according to claim 17, wherein each of L_1 and L_2 represents a fluorescence donor ~~or~~ or a fluorescence quencher ~~pair~~.

19. (currently amended) The compound according to claim 18, wherein each of L_1 and L_2 represents a donor or an acceptor in a FRET pair.

20. (previously presented) The compound according to claim 15, wherein L_2 is avidin, streptavidin or biotin.

21. (previously presented) The compound according to claim 15, wherein L_2 is a moiety covalently attached to a solid support.

22. (currently amended) The compound according to claim 15, wherein L_2 is a linear poly(arginine) of D- and/or L-arginine with 6-15

arginine residues, an oligomer of 6-50 subunits wherein at least one subunit has an attached guanidinium group ~~a linear polymer of 6-15 subunits each carrying a guanidinium group, oligomers or short length polymers of 6-50 subunits, a portion of which have attached guanidinium groups, or parts~~ a partial amino acid sequence of a ~~sequence of the~~ HIV-tat protein.

23-43 (cancelled)

44. (currently amended) A method for detecting ~~and/or manipulating~~ a protein of interest, wherein the protein of interest is fused to an AGT, the AGT fusion protein is contacted with a compound of formula (1) according to claim 1, and the AGT fusion protein is detected ~~and optionally further manipulated~~ using the label L₁ and/or L₂ in a system designed for recognizing and/or handling the label.

45. (currently amended) The method according to claim 44, wherein in the compound of formula (1) ~~label~~ L₂ is a solid support, and the AGT fusion protein contacted with the compound of formula (1) is separated from the compound of formula (1) by filtration or centrifugation or separation of magnetic beads.

46. (currently amended) The method according to claim 44, wherein in the compound of formula (1) ~~label~~ L₁ is one member and ~~label~~ L₂ the other member of two interacting ~~spectroscopic probes~~ chromophores or fluorophores L₁ / L₂, wherein energy can be transferred nonradiatively through dynamic or static quenching, and the AGT fusion protein is detected by fluorescence.

47. (currently amended) The method according to claim 44 for detecting ~~and/or manipulating~~ a protein of interest, wherein the protein of interest is fused with a mutant AGT, the mutant AGT fusion protein is contacted with a mixture of

(a) a compound of formula (1) wherein R_1 ~~is a group~~ additionally comprises R_5 , wherein R_5 is a substituted or unsubstituted cycloalkyl, cycloalkenyl or heterocyclyl group and which does not react with the mutant AGT, and

(b) another compound of formula (1), which reacts with the mutant AGT fusion protein, and the mutant AGT fusion protein is detected ~~and optionally further manipulated~~ using the label in a system designed for recognizing and/or handling the label.